



University of Groningen

Physical frailty in late-life depression: evidence for a depression-frailty subtype?

Arts, Matheus

DOI:

[10.33612/diss.147370083](https://doi.org/10.33612/diss.147370083)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Arts, M. (2020). *Physical frailty in late-life depression: evidence for a depression-frailty subtype?*. University of Groningen. <https://doi.org/10.33612/diss.147370083>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter I

General introduction

Due to the baby boom after World War II and an (still) increasing life expectancy, around 25% of the Dutch population is 65 years of older, of which more than 30% is older than 80 years in 2050 (www.cbs.nl). One of the most challenging aspects of an ageing population is to preserve or improve a good physical and mental health of older adults (WHO, 2015). From a medical perspective, interventions should be focused on those people with the highest risk of adverse health outcomes (Fried et al., 2001; Collard et al., 2012; WHO, 2015). To this end, frailty has been introduced as a core concept in geriatric medicine as frail older persons have the highest risks of adverse health outcomes and are high users of community resources, hospitalization, and nursing homes (Morley et al., 2013). While the relevance of frailty is increasingly recognized in geriatric as well as general medicine, this concept is largely ignored in geriatric psychiatry. This is remarkable, as several psychiatric disorders, in particular depression, negatively interfere with life-expectancy (Cuijpers et al., 2014; Brandao et al., 2019). This thesis focusses on the validity and clinical relevance of frailty in late-life depression.

Depression - a disorder of accelerated aging

Globally, depression contributes significantly to the burden of disease (Lépine & Briley, 2011) and is identified as an independent predictor for the onset of many chronic somatic diseases (Penninx et al., 2013). Current predictions indicate that by 2020, depression will be the second leading cause of disease burden, and by 2030 it probably will be the leading cause worldwide (www.who.int). In later life, depression has higher recurrence rates, a more chronic course, and a higher level of co-morbidity with cognitive impairment and somatic diseases (Comijs et al., 2011; Schaakxs et al., 2018). The prevalence of depressive disorder according to the diagnostic criteria of the DSM or ICD classification systems is estimated at 1.8% among community-dwelling people aged 55 years and older (Beekman et al, 1999) and 7.2% among older adults aged 75 years and over (independent of the living

arrangement) (Luppa et al., 2012). Furthermore, in older populations, subthreshold depression is more prevalent than a major depressive disorder (Hasin et al., 2005; Fiske et al., 2009), with prevalence rates estimated between 13.5% and 17.5% (Beekman et al., 1999; Luppa et al., 2012). As these latter estimates are often based on self-report symptom scales, confounding by underlying and/or subthreshold somatic diseases cannot be excluded (Thombs et al., 2010). Thus far, studies on the impact of depression on the onset of somatic diseases have not included physical frailty as a confounding and/or mediating variable (Collard et al., 2015).

The hypothesis that depression itself is a clinical condition of accelerated aging (Verhoeven et al., 2014) has originated from findings that depression is prospectively associated with the onset of chronic somatic diseases and an increased mortality rate independent of lifestyle characteristics (Cuijpers et al., 2013; Penninx et al., 2013). This hypothesis is further fed by associations between ageing-related biomarkers and depressive disorder (e.g. Howren et al., 2009; Dowlati et al., 2010; Wolkowitz et al., 2011; Pan et al., 2012; Verhoeven et al., 2014; Morrison et al., 2019). Most studies on the association between ageing biomarkers and depression did not take frailty into account. This might be relevant, as frailty and depression share partly overlapping diagnostic criteria as well as assumed underlying pathophysiological mechanisms (Mezuk et al., 2012), see below for more detailed discussion. Therefore, in this thesis, we examined the association between frailty and ageing-related biomarkers of ageing among depressed older patients. We examined specifically low graded inflammation (**chapter 2**), shortened telomere length (**chapter 3**), and lowered vitamin D levels (**chapter 4**).

Low graded inflammation - This process of accelerated aging is thought to occur as a result of dysregulations in immune responses that may contribute to changes of the affective and cognitive neural systems associated with the development of late-

life depression (Alexopoulos & Morimoto, 2011). The aging process creates a condition of chronic low-grade inflammation, also called “inflamm-ageing” (Grolleau-Julius et al., 2010) that leads to the production of several pro-inflammatory cytokines (Alexopoulos & Morimoto, 2011). Many studies focused on the association of cytokines and depression, and found that C-reactive protein as well as pro-inflammatory cytokine levels were increased in older adults with depressive symptoms (Dentino et al., 1999; Penninx et al., 2003; Tiemeier et al., 2003; Bremner et al., 2008; Milaneschi et al., 2009; Zalli et al., 2016), albeit negative studies have also been reported (Stewart et al., 2009; Matsushima et al., 2015).

Shortened telomere length - Telomere length is widely considered as a marker of cellular aging, as shortened telomeres in white blood cells are associated with increased mortality rates (Cawthon et al., 2003; Honig et al., 2006) and the onset of various age-related diseases (Collado et al., 2007; Willeit et al., 2010). Associations between shortened leucocyte telomere length and depression in young- and middle-aged patients have been found (e.g. Epel et al., 2004; Damjanovic et al., 2007; Kananen et al., 2010; Tyrka et al., 2010; Garcia-Rizo et al., 2013; Verhoeven et al., 2014; Lin et al., 2016). However, these findings could not be replicated in the Netherlands Study of Depression in Older persons (NESDO) (see **appendix**) (Schaakxs et al., 2015). This has amongst others been explained by the fact that late-life depression has a more heterogeneous nature as compared to depression earlier in life, which may mask (small) effects pertaining to specific subgroups. Whether this subgroup is characterised by the presence of physical frailty has not been examined yet.

Vitamin D deficiency - Vitamin D is involved in various cellular ageing processes, including proliferation, cellular differentiation, and apoptosis (Pusceddu et al., 2015; Berridge, 2017). In human studies, an inverse association between vitamin D and

ageing-related diseases and mortality has been reported (Pusceddu et al., 2015). A growing body of evidence relates lower vitamin D levels to physical frailty (Shardell et al., 2009; Chang et al., 2010; Wilhelm-Leen et al., 2010; Ensrud et al., 2011; Smit et al., 2012; Hirani et al., 2013; Pabst et al., 2015; Ju et al., 2018). Low vitamin D levels may to some extent predict frailty and could represent a biomarker for identifying older adults at risk for frailty (Buta et al., 2017). Underlying mechanisms of this relationship may be the involvement of vitamin D in sarcopenia, which is closely related to frailty, as vitamin D stimulates Ca^{2+} and phosphorus transport, muscle contraction, and muscle differentiation (Garcia et al., 2011; Ju et al., 2018). The association between vitamin D deficiency and depression has been widely debated (Kerr et al., 2015). While low levels of vitamin D are consistently associated with depression (Anglin et al., 2013; Spedding, 2014), meta-analyses of treatment studies have reported no significant effects on depression after vitamin D supplementation (Li et al., 2013; Shaffer et al., 2014).

Frailty - a concept of biological aging

Frailty is a vulnerability state characterized by poor resolution of homeostasis after a stressor, placing persons at risk of iatrogenic damage, dependency and death (Morley et al., 2013). A consensus group, consisting of frailty-experts and delegates from six major international societies, defined *physical* frailty as an important medical syndrome with multiple causes and contributors that is characterized by diminished strength, endurance, and reduced physiologic function (Morley et al., 2013). This concept of physical frailty was considered relevant to target geriatric care to the most vulnerable patients. Promising prevention and treatment strategies for physical frailty included reduction or prevention of polypharmacy, more physical exercise, protein-calorie supplementation, and regular monitoring of vitamin D status to provide tailored care (Morley et al., 2013). Among community-dwelling

persons aged 65 years and over, the mean prevalence rate of physical frailty was 9.9% (range 4.0 – 17.0%) (Collard et al., 2012).

Two dominant operationalisations are 1) the deficit accumulation model (Frailty Index), stating that the proportion of ageing-related deficits reflects biological age on top of chronological age, and 2) the Fried Frailty Phenotype, which mark an underlying physiological state of multisystem and energy dysregulation (Fried et al., 2001; Mitnitski et al., 2001; Cesari et al., 2014). According to Cesari et al (2014), the Fried Frailty Phenotype and the Frailty Index should not be considered substitutable with each other, but as complementary constructs.

The Frailty Index has been developed in the Canadian Study of Health and Aging (CSHA), a five-year prospective cohort study which included 10,263 older adults aged 65 years and older (Rockwood et al., 2005). The Frailty Index was originally calculated by measuring the absence or presence of 92 clinical deficits (Mitnitski et al., 2001; Mitnitski et al., 2002; Rockwood et al., 2005). Ongoing research has shown that the Frailty Index is not dependent on the number of deficits taken into account, as long as at least 30 deficits from several health domains have been included. However, the Frailty Index may substantially overlap with psychiatric conditions, as several measures of mental and cognitive capacity are included.

The Fried Frailty Phenotype is defined as the presence of at least three out of five criteria, i.e. weight loss, exhaustion, weakness, slowness, and reduced physical activity. This definition has been validated in the Cardiovascular Health Study (CHS), showing predictive value for incident falls, worsening mobility, hospitalization, and death during follow-up independent of multimorbidity or disability (Fried et al., 2001). Although the Fried Frailty Phenotype has been used by many researchers, the operationalisation of the specific components slightly differs between studies. Some

studies have shown that self-report indices of the five components also provide a reliable proxy of the Fried Frailty Phenotype (Lee et al., 2017; Papachristou et al., 2017). In this thesis, frailty has been defined according to the Fried Frailty Phenotype (Fried et al., 2001). We assumed that this model would be most appropriate for application in a psychiatric setting in order to prevent symptom overlap and confounding with psychiatric disorders. It allows us to study frailty in relation to psychiatric disorders without including psychopathology in the broad definition of frailty (Fried et al., 2001; Collard et al., 2012). Hereby, we would be able to disentangle the relationship between psychopathology, frailty, and underlying ageing mechanisms as well as the clinical impact of frailty in geriatric psychiatry (Collard et al., 2012).

Frailty and depression - approaching concepts with increasing age?

Depression is, analogously to frailty, associated with negative health outcomes, including somatic diseases and increased mortality rates (Penninx et al, 2013; Cuijpers et al, 2014). Nonetheless, in the hallmark study of Fried and colleagues (2001) on the Fried Frailty Phenotype, depressed patients or persons on antidepressants were a priori excluded to prevent being classified as physically frailty due to the presence of only one (psychiatric) disorder (Fried et al, 2001). This might unnecessarily have contributed to the neglect of the concept of frailty in geriatric psychiatry.

A narrative review on the association between depression and frailty points not only to overlap between both conditions, but also to similarities in the determinants and consequences of both concepts (Mezuk et al., 2012). The overlap between both constructs has been confirmed in empirical studies. Using latent class analyses among adults aged 41 – 96 years, depression and physical frailty appeared to be distinct concepts, although 100% of the severely depressed persons were classified

as physically frail (Mezuk et al., 2013). Moreover, factor analyses on the criteria for depression and frailty in the Health and Retirement Study, showed that depression and frailty represent two distinct, but highly associated dimensions (Lohman et al., 2016). These findings suggest that frailty and depression in later life are difficult to disentangle. A recent meta-analysis on the association between depression and frailty among older adults estimated that across cross-sectional studies, 40.4% of depressed persons were frail and that 38.6% of frail persons were depressed (Soysal et al., 2017). A limited number of longitudinal studies confirmed these findings (Soysal et al., 2017). Therefore, frailty and late-life depression may partly share some of their etiological pathways as well as their phenotypic expression (Ni Mhaolain et al., 2012). Nonetheless, only three studies included in the above mentioned meta-analysis have assessed depression according to the diagnostic criteria of the DSM, among which a study from the Netherlands Study of Depression in Older persons (NESDO). In NESDO, only 27.2% of the patients suffering from a DSM-IV defined depressive disorder were classified physically frail, which could not be explained by overlapping criteria for frailty and depression (Collard et al., 2014). These prevalence rates are comparable to the prevalence of frailty among populations with chronic somatic conditions in which the clinical relevance of frailty becomes more and more evident (Denfeld et al., 2017).

Thus far, frailty has neither been examined as a determinant of the course of depression nor of an increased mortality rate in late life depression, as done in **chapter 7** and **chapter 8**.

Frailty and common (neuro)psychiatric symptoms in later life

While studies on the association between psychopathology and frailty primarily focusses on depression, frailty is probably as relevant for other (neuro)psychiatric disorders (Andrew & Rockwood, 2007). Therefore, in this thesis we also investigate the association between physical frailty and neuropsychiatric disorders in later life.

We will focus on “cognitive impairment” and “medically unexplained symptoms” (MUS).

Cognitive impairment - The association between physical frailty and cognitive performance is complex (Malmstrom et al., 2013; Panza et al., 2014). A systematic review has identified eleven longitudinal studies that show the predictive value of physical frailty for subsequent cognitive decline or dementia (Robertson et al., 2013). Although less often studied, cognitive impairment conversely may be a risk factor of physical frailty (Robertson et al., 2013; Nishiguchi et al., 2015). Some researchers have included cognitive deficits as a component of physical frailty for two reasons. Firstly, adding cognitive performance to a frailty index adds to its predictive validity for adverse health outcomes (Ávila-Funes et al., 2009). Secondly, significant overlap exists in the mechanisms underlying physical frailty and cognitive impairment (Halil et al., 2015). In 2013, the concept “cognitive frailty” has been introduced by a consensus panel to emphasize the important role of brain aging (Kelaiditi et al., 2013). Cognitive frailty was defined as a subtype of physical frailty and characterized by the concurrence of cognitive deficits in physically frail older persons in the absence of an accompanying neurological disorder (Kelaiditi et al., 2013). Cognitive frailty may represent a prodromal phase for neurodegenerative diseases (Kelaiditi et al., 2013), but its potential for reversibility makes it an ultimate target for early intervention (Kelaiditi et al., 2013). Late-life depression can also have a detrimental effect on cognitive performance (Panza et al., 2009; Wilkins et al., 2009; Korten et al., 2014; Panza et al., 2014). Nonetheless, the association between physical frailty and cognitive performance has been hardly studied among clinically depressed older persons (as will be done in **chapter 5**).

Medically unexplained symptoms - Medically unexplained symptoms (MUS) are defined as physical symptoms that, after appropriate medical assessment, cannot

be explained in terms of a conventionally defined medical disease (Wessely, 1999; Beck, 2008). Patients with medically unexplained symptoms (MUS) often describe a low quality of life and frequently suffer from co-morbid anxiety and depressive disorders (De Waal et al., 2004). This gives rise to high levels of health care consumption in the search for an organic origin of complaints and places especially older persons at risk for iatrogenesis (Smith et al., 2005). Besides a small pilot study (Benraad et al., 2013), no studies have been conducted on the physical performance of older patients with MUS. This pilot study suggests that MUS in later life might be related to the presence of frailty. Therefore, in this thesis we want to examine the association between frailty and somatic comorbidity among older patients with MUS and older patients with Medically Explained Symptoms (MES) who have participated in the “Older Persons with medically Unexplained Symptoms” (OPUS) study (see **appendix**). We will particularly explore the association between the severity of MUS and frailty (**chapter 6**).

Aims and outlines of this thesis

The main aim of this thesis is to study physical frailty in an older population of depressed older adults. This thesis consists of three parts.

Part I:

Ageing-related biomarkers of physical frailty in late-life depression.

The huge variability in biological ageing among individuals who live in a population characterised by an increasing life expectancy has stimulated the search for biomarkers of biological ageing. Ageing-related biomarkers should be associated with physical frailty as a clinical phenotype of ageing, but also with conditions of accelerated ageing like depression. This first part of the thesis focuses on the relationship between physical frailty and age-related biomarkers in a population of depressed older adults. As physical frailty and late-life depression partly overlap and the association between depression and ageing-related biomarkers decreases with age, we examined whether associations between ageing-related biomarkers and physical frailty were moderated by the presence of a late-life depression within the Netherlands Study of Depression in Older persons (NESDO). The specific research questions that have to be answered are (numbers corresponding with chapters):

- **Chapter 2:** Is physical frailty associated with low-grade inflammation in a cohort of depressed older patients?
- **Chapter 3:** Is physical frailty associated with telomere length in a cohort of depressed older patients?
- **Chapter 4:** Is physical frailty associated with vitamin D levels in a cohort of depressed older patients?

Part II:

Association between physical frailty and common (neuro)psychiatric symptoms.

The operationalisation of the concept of (physical) frailty is still debated and psychiatric disorders beyond depression have also been associated with increased mortality rates. Therefore, the idea underlying this second part is the question whether the physical frailty phenotype should be broadened with neuropsychiatric symptoms beyond depression. Firstly, we set up a study to explore whether physical frailty is associated with poorer cognitive functioning in a sample of depressed older adults. Secondly, we examined the level of physical frailty and somatic comorbidity in older patients with medically unexplained symptoms (MUS) and compared this to patients with medically explained symptoms (MES). The following research questions have to be answered:

- **Chapter 5:** Is cognitive impairment in late-life depression associated with physical frailty?
- **Chapter 6:** Are medically unexplained symptoms in later life associated with physical frailty?

Part III:

Adverse health outcomes of physical frailty in late-life depression.

The third part of the thesis is more clinically oriented by focusing on the predictive value of physical frailty in late-life depression. First, we investigated the prognostic impact of frailty on the outcome and course of late-life depression. Secondly, we examined whether physical frailty predicts mortality among clinically depressed older persons. The corresponding research questions are:

- **Chapter 7:** Does physical frailty negatively impacts the course of depression in later life?
- **Chapter 8:** Does physical frailty increase mortality risk in late-life depression?

Appendix

Netherlands Study of Depression in Older persons (NESDO)

The NESDO study was set up to study neurobiological, physical, and psychosocial determinants and the course of late-life depression. NESDO is a multisite, naturalistic cohort study (Comijs et al., 2011) and included 378 older (≥ 60 years) participants with a 6-months diagnosis of major or minor depression or dysthymia, as well as a comparison group of 132 never depressed participants (≥ 60 years). Depressed participants were recruited in primary care as well as secondary and tertiary mental health care institutes. Non-depressed controls were recruited from general practices only. Exclusion criteria were a suspected or established diagnosis of dementia or a Mini Mental State Examination-score <18 (Folstein et al., 1975) and insufficient command of the Dutch language. Participants were followed-up for six years including self-report symptom severity scales every six months by post and full re-examinations at two- and six-year follow-up (site visits). A more detailed description of the NESDO study can be found elsewhere (Comijs et al., 2011; Comijs et al., 2015; Jeuring et al., 2018).

Older Persons with medically Unexplained Symptoms (OPUS) Study

The OPUS study was set up to explore the biopsychosocial determinants and course of medically unexplained symptoms (MUS) in later life. A mixed method design has been adopted by including both quantitative and qualitative data collection. In this thesis, only the quantitative baseline assessment will be used, which consists of a case-control study of 118 older MUS-patients (>60 years) with chronic MUS (>3 months) to 154 older patients with chronic (>3 months) medically explained symptoms (MES), such as diabetes or COPD. To compose a sample of patients representing the whole severity range of MUS, participants with MUS and MES were recruited in the community (advertisements in local newspapers), in primary care, and in secondary health care (outpatient mental health clinic as well as outpatient geriatric

department of a University Medical Center). Exclusion criteria were the presence of a primary psychotic disorder, cognitive impairment, terminal illness, severe auditory and/or visual limitations, and insufficient command of the Dutch language.

References

Alexopoulos GS, Morimoto SS. The inflammation hypothesis in geriatric depression. *Int J Geriatr Psychiatry* 2011;26(11):1109-1118.

Andrew MK, Rockwood K. Psychiatric illness in relation to frailty in community-dwelling elderly people without dementia: a report from the Canadian study of health and aging. *Can J Aging* 2007;26(1):33-38.

Anglin RE, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *Br J Psychiatry* 2013;202:100-7.

Avila-Funes JA, Amieva H, Barberger-Gateau P, Le Goff M, Raoux N, Ritchie K, Carrière I, Tavernier B, Tzourio C, Gutiérrez-Robledo LM, Dartiques JF. Cognitive impairment improves the predictive validity of the phenotype of frailty for adverse health outcomes: The three-city study. *J Am Geriatr Soc* 2009;57(3):453-461.

Beck EJ. A developmental perspective on functional somatic symptoms. *J Pediatr Psychol* 2008;33(5):547-562.

Beekman AT, Copeland JR, Prince MJ. Review of community prevalence of depression in later life. *Br J Psychiatry* 1999;174:307-311.

Benraad CE, Hilderink PH, van Driel DT, Disselhorst LG, Lubberink B, van Wolferen L, Olde Rikkert MG, Oude Voshaar RC. Physical functioning in older persons with somatoform disorders: a pilot study. *J Am Dir Assoc* 2013;14(1):75e9-13.

Berridge MJ. Vitamin D deficiency accelerates ageing and age-related diseases: a novel hypothesis. *J Physiol* 2017;595(22):6825–6836.

Brandao DJ, Fontenelle LF, da Silva SA, Menezes PR, Pastor-Valero M. Depression and excess mortality in the elderly living in low- and middle-income countries: systematic review and meta-analysis. *Int J Geriatr Psychiatry* 2019;34(1):22-30.

Bremmer MA, Beekman AT, Deeg DJ, Penninx BW, Dik MG, Hack CE, Hoogendijk WJ. Inflammatory markers in late-life depression: results from a population-based study. *J Affect Disord* 2008;106(3):249-255.

Buta B, Choudhury PP, Xue Q, Chaves P, Bandeen-Roche K, Shardell M, Semba RD, Walston J, Michos ED, Appel LJ, McAdams-DeMarco M, Gross A, Yasar S, Ferrucci L, Fried LP, Kalyani RR. The association of vitamin D deficiency and incident frailty in older women: the role of cardiometabolic diseases. *J Am Geriatr Soc* 2017;65(3):619-624.

Cawthon RM, Smith KR, O'Brien E, Sivatchenko A, Kerber RA. Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet* 2003;361(9355):393–395.

Centraal bureau voor de Statistiek (CBS). StatLine: prognose bevolking, geslacht en leeftijd. 2019. (www.cbs.nl)

Cesari M, Gambassi G, van Kan GA, Vellas B. The frailty phenotype and the frailty index: different instruments for different purposes. *Age Ageing* 2014;43(1):10-2.

Chang CI, Chan DC, Kuo KN, Hsiung CA, Chen CY. Vitamin D insufficiency and frailty syndrome in older adults living in a Northern Taiwan community. *Arch Gerontol Geriatr* 2010;50(1):S17-21.

Collado M, Blasco MA, Serrano M. Cellular senescence in cancer and aging. *Cell* 2007;130:223–233.

Collard RM, Oude Voshaar RC. Frailty; een kwetsbaar begrip. *Tijdschrift voor Psychiatrie* 2012;54(1):59-69.

Collard RM, Comijs HC, Naarding P, Oude Voshaar RC. Physical frailty: vulnerability of patients suffering from late-life depression. *Aging Ment Health* 2014;18(5):570-578.

Collard RM, Arts M, Comijs HC, Naarding P, Verhaak PFM, de Waal MW, Oude Voshaar RC. The role of frailty in the association between depression and somatic comorbidity: Results from baseline data of an ongoing prospective cohort study. *Int J Nurs Studies* 2015;16:509-514.

Comijs HC, Van Marwijk HW, Van Der Mast RC, Naarding P, Oude Voshaar RC, Beekman AT, Boshuisen M, Dekker J, Kok R, de Waal MW, Penninx BW, Stek ML, Smit JH. The Netherlands study of depression in older persons (NESDO); a prospective cohort study. *BMC Res Notes* 2011;4:524.

Comijs HC, Nieuwesteeg J, Kok R, van Marwijk HW, van der Mast RC, Naarding P, Oude Voshaar RC, Verhaak P, de Waal MW, Stek ML. The two-year course of late-life depression: results from the Netherlands study of depression in older persons. *BMC Psychiatry* 2015;15:20.

Cuijpers P, Vogelzangs N, Twisk J, Kleiboer A, Li J, Penninx BW. Differential mortality rates in major and subthreshold depression: meta-analysis of studies that measured both. *Br J Psychiatry* 2013;202:22-27.

Cuijpers P, Vogelzangs N, Twisk J, Kleiboer A, Li J, Penninx BW. Comprehensive meta-analysis of excess mortality in depression in the general community versus patients with specific illnesses. *Am J Psychiatry* 2014;171(4):453-462.

Damjanovic AK, Yang Y, Glaser R, Kiecolt-Glaser JK, Nguyen H, Laskowski B, Zou Y, Beversdorf DQ, Weng NP. Accelerated telomere erosion is associated with a declining immune function of caregivers of Alzheimer's disease patients. *J Immunol* 2007;179:4249–4254.

De Waal MW, Arnold IA, Eekhof JA, van Hemert AM. Somatoform disorders in general practice: prevalence, functional impairment and comorbidity with anxiety and depressive disorders. *Br J Psychiatry* 2004;184:470-476.

Denfeld QE, Winters-Stone K, Mudd JO, Gelow JM, Kurdi S, Lee CS. The prevalence of frailty in heart failure: a systematic review and meta-analysis. *Int J Cardiol* 2017;236:283-289.

Dentino AN, Pieper CF, Rao MK, Currie MS, Harris T, Blazer DG, Cohen HJ. Association of interleukin-6 and other biologic variables with depression in older people living in the community. *J Am Geriatr Soc* 1999;47(1):6-11.

Dowlati Y, Herrmann N, Swardfager W, Liu H, Sham L, Reim EK, Lanctôt KL. A Meta-Analysis of Cytokines in Major Depression. *Biol Psychiat* 2010;67(5):446-457.

Ensrud KE, Blackwell TL, Cauley JA, Cummings SR, Barrett-Connor E, Dam TT, Hoffman AR, Shikany JM, Lane NE, Stefanick ML, Orwoll ES, Cawthon PM, Osteoporotic Fractures in Men Study Group. Circulating 25-hydroxyvitamin D levels

and frailty in older men: the osteoporotic fractures in men study. *J Am Geriatr Soc* 2011;59(1):101-6.

Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, Cawthon RM. Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci USA* 2004;101:1173–1182.

Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annu Rev Clin Psychol* 2009;5:363-389.

Folstein MF, Folstein SE, McHugh PR. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12(3):189-1898.

Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, Seeman T, Tracy R, Kop WJ, Burke G, McBurnie MA, Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56(3):M146-156.

Garcia LA, King KK, Ferrini MG, Norris KC, Artaza JN. 1,25(OH)₂vitamin D₃ stimulates myogenic differentiation by inhibiting cell proliferation and modulating the expression of promyogenic growth factors and myostatin in C2C12 skeletal muscle cells. *Endocrinology* 2011;152(8):2976–2986.

Garcia-Rizo C, Fernandez-Egea E, Miller BJ, Oliveira C, Justicia A, Griffith JK, Heaphy CM, Bernadro M, Kirkpatrick B. Abnormal glucose tolerance, white blood cell count, and telomere length in newly diagnosed, antidepressant-naïve patients with depression. *Brain Behav Immun* 2013;28:49-53.

Grolleau-Julius A, Ray D, Yung RL. The role of epigenetics in aging and autoimmunity. *Clin Rev Allergy Immunol* 2010;39:42-50.

Halil M, Cemal Kizilarlanoglu M, Emin Kuyumcu M, Yesil Y, Cruz Jentoft AJ. Cognitive aspects of frailty: Mechanisms behind the link between frailty and cognitive impairment. *J Nutr Health Aging* 2015;19(3):276-283.

Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Arch Gen Psychiatry* 2005;62:1097-1106.

Hirani V, Naganathan V, Cumming RG, Blyth F, Le Couteur DG, Handelsman DJ, Waite LM, Seibel MJ. Associations between frailty and serum 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D concentrations in older Australian men: the Concord Health and Ageing in Men Project. *J Gerontol A Biol Sci Med Sci* 2013;68(9):1112-21.

Honig LS, Schupf N, Lee JH, Tang MX, Mayeux R. Shorter telomeres are associated with mortality in those with APOE epsilon4 and dementia. *Ann Neurol* 2006;60(2):181–187.

Howren MB, Lamkin DM, Suls J. Associations of Depression With C-Reactive Protein, IL-1, and IL-6: A Meta-Analysis. *Psychosom Med* 2009;71(2):171-186.

Jeuring HW, Stek ML, Huisman M, Oude Voshaar RC, Naarding P, Collard RM, van der Mast RC, Kok RM, Beekman ATF, Comijs HC. A six-year prospective study of the prognosis and predictors in patients with late-life depression. *Am J Geriatr Psychiatry* 2018;26(9):985-997.

Ju SY, Lee JY, Kim DH. Low 25-hydroxyvitamin D levels and the risk of frailty syndrome: a systematic review and dose-response meta-analysis. *BMC Geriatr* 2018;18(1):206.

Kananen L, Surakka I, Pirkola S, Suvisaari J, Lonqvist J, Peltonen L, Ripatti S, Hovatta I. Childhood adversities are associated with shorter telomere length at adult age both in individuals with an anxiety disorder and controls. *PloS One* 2010;5(5):e10826.

Kelaiditi E, Cesari M, Canevelli M, van Kan GA, Ousset PJ, Gillette-Guyonnet S, Ritz P, Duveau F, Soto ME, Provencher V, Nourhashemi F, Salvá A, Robert P, Andrieu S, Rolland Y, Touchon J, Fitten JL, Vellas B; IANA/IAGG. Cognitive frailty: Rational and definition from an (I.A.N.A./I.A.G.G.) international consensus group. *J Nutr Health Aging* 2013;17(9):726-734.

Kerr DCR, Zava DT, Piper WT, Saturn SR, Frei B, Gombart AF. Associations between vitamin D levels and depressive symptoms in healthy young adult women. *Psychiatry Res* 2015;227(1):46-51.

Korten NC, Penninx BW, Kok RM, Stek ML, Oude Voshaar RC, Deeg DJ, Comijs HC. Heterogeneity of late-life depression: Relationship with cognitive functioning. *Int Psychogeriatr* 2014;26(6):953-963.

Lee L, Patel T, Costa A, Bryce E, Hillier LM, Slonim K, Hunter SW, Heckman G, Molnar F. Screening for frailty in primary care: accuracy of gait speed and handgrip strength. *Can Fam Physician* 2017;63(1):e51-e57.

Lépine JP, Briley M. The increasing burden of depression. *Neuropsychiatr Dis Treat* 2011;7(1):3-7.

Li L, Wu B, Liu JY, Yang LB. Vitamin D receptor gene polymorphisms and type 2 diabetes: a meta-analysis. *Arch Med Res* 2013;44(3):235-241.

Lin PY, Huang YC, Hung CF. Shortened telomere length in patients with depression: a meta-analytic study. *J Psychiatr Res* 2016;76:84-93.

Lohman M, Dumenci L, Mezuk B. Depression and frailty in later life: evidence for a common vulnerability. *J Gerontol B Psychol Sci Soc Sci* 2016;71(4):630-640.

Luppa M, Sikorski C, Luck T, Weyerer S, Villringer A, König HH, Riedel-Heller SG. Prevalence and risk factors of depressive symptoms in latest life-results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). *Int J Geriatr Psychiatry* 2012;27(3):286-295.

Malmstrom TK, Morley JE. Frailty and cognition: Linking two common syndromes in older persons. *J Nutr Health Aging* 2013;17(9):723-725.

Matsushima J, Kawashima T, Nabeta H, Imamura Y, Watanabe I, Mizoguchi Y, Kojima N, Yamada S, Monji A. Association of inflammatory biomarkers with depressive symptoms and cognitive decline in a community-dwelling healthy older sample: a 3-year follow-up study. *J Affect Disord.* 2015;173:9-14.

Mezuk B, Edwards L, Lohman M, Choi M, Lapane KL. Depression and frailty in later life: a synthetic review. *Int J Geriatr Psychiatry* 2012;27(9):879-892.

Mezuk B, Lohman M, Dumenci L, Lapane KL. Are depression and frailty overlapping syndromes in mid- and late-life? A latent variable analysis. *Am J Geriatr Psychiatry* 2013;21(6):560-9.

Milaneschi Y, Corsi AM, Penninx BW, Bandinelli S, Guralnik JM, Ferrucci L. Interleukin-1 receptor antagonist and incident depressive symptoms over 6 years in older persons: the InCHIANTI study. *Biol Psychiatry* 2009;65(11):973-978.

Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *ScientificWorldJournal* 2001;1:323-336.

Mitnitski AB, Graham JE, Mogilner AJ, Rockwood K. Frailty, fitness and late-life mortality in relation to chronological and biological age. *BCM Geriatr* 2002;2:1.

Morley JE, Vellas B, Abellan van Kan G, Anker SD, Bauer JM, Bernabei R, Cesari M, Chumlea WC, Doehner W, Evans J, Fried LP, Guralnik JM, Katz PR, Malmstrom TK, McCarter RJ, Gutierrez Robledo LM, Rockwood K, von Haehling S, Vandewoude MF, Walston J. Frailty consensus: a call to action. *J Am Dir Assoc* 2013;14(6):392-397.

Morrison FG, Logue MW, Guetta R, Maniates H, Stone A, Schichman SA, McGlinchey RE, Milberg WP, Miller MW, Wolf EJ. Investigation of bidirectional longitudinal associations between advanced epigenetic age and peripheral biomarkers of inflammation and metabolic syndrome. *Aging (Albany NY)* 2019;11(11):3487-3504.

Ni Mhaoláin AM, Fan CW, Romero-Ortuno R, Cogan L, Cunningham C, Kenny RA, Lawlor B. Frailty, depression, and anxiety in later life. *Int Psychogeriatr* 2012;24(8):1265-1274.

Nishiguchi S, Yamada M, Fukutani N, Adachi D, Tashiro Y, Hotta T, Morino S, Shirooka H, Nozaki Y, Hirata H, Yamaguchi M, Arai H, Tsuboyama T, Aoyama T. Differential association of frailty with cognitive decline and sarcopenia in community-dwelling older adults. *J Am Med Dir Assoc* 2015;16(2):120-124.

Pabst G, Zimmermann AK, Huth C, Koenig W, Ludwig T, Zierer A, Peters A, Thorand B. Association of low 25-hydroxyvitamin D levels with the frailty syndrome in an aged population: results from the KORA-age Augsburg study. *J Nutr Health Aging* 2015;19(3):258-264.

Pan A, Keum N, Okereke OI, Sun Q, Kivimaki M, Rubin RR, Hu FB. Bidirectional association between depression and metabolic syndrome: a systematic review and meta-analysis of epidemiological studies. *Diabetes Care* 2012;35(5):1171-1180.

Panza F, D'Introno A, Colacicco AM, Capurso C, Del Parigi A, Caselli RJ, Frisardi V, Scapicchio P, Chiloio R, Scafato E, Gandin C, Vendemiale G, Capurso A, Solfrizzi V. Temporal relationship between depressive symptoms and cognitive impairment: The Italian Longitudinal Study on Aging. *J Alzheimers Dis* 2009;17(4):899-911.

Panza F, Solfrizzi V, Tortelli R, Resta F, Sabbá C, Logroscino G. Prevention of late-life cognitive disorders: diet-related factors, dietary patterns, and frailty models. *Curr Nutr Rep* 2014;3:110e129.

Papachristou E, Wannamethee SG, Lennon LT, Papacosta O, Whincup PH, Iliffe S, Ramsay SE. Ability of self-reported frailty components to predict incident disability, falls, and all-cause mortality: results from a population-based study of older British men. *J Am Med Dir Assoc* 2017;18(2):152-157.

Penninx BW, Kritchevsky SB, Yaffe K, Newman AB, Simonsick EM, Rubin S, Ferrucci L, Harris T, Pahor M. Inflammatory markers and depressed mood in older persons: results from the Health, Aging and Body Composition study. *Biol Psychiatry* 2003;54(5):566-572.

Penninx BWJH, Milaneschi Y, Lamers F, Vogelzangs N. Understanding the somatic consequences of depression: biological mechanisms and the role of depression symptom profile. *BMC Medicine* 2013;11:129.

Pusceddu I, Farrell CL, Di Pierro AM, Jani E, Herrmann W, Herrmann M. The association of vitamin D deficiency and incident frailty in older women: the role of cardiometabolic diseases. *Clin Chem Lab Med* 2015;53(11):1661–1678.

Robertson DA, Savva GM, Kenny RA. Frailty and cognitive impairment - a review of the evidence and causal mechanisms. *Ageing Res Rev* 2013;12:840-851.

Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489-495.

Schaakxs R, Verhoeven JE, Oude Voshaar RC, Comijs HC, Penninx BW. Leukocyte telomere length and late-life depression. *Am J Geriatr Psychiatry* 2015;23(4):423-432.

Schaakxs R, Comijs HC, Lamers F, Kok RM, Beekman ATF, Penninx BJWH. Associations between age and the course of major depressive disorder: a 2-year longitudinal cohort study. *Lancet Psychiatry* 2018;5(7):581-590.

Shaffer JA, Edmondson D, Wasson LT, Falzon L, Homma K, Ezeokoli N, Li P, Davidson KW. Vitamin D supplementation for depressive symptoms: a systemic review and meta-analysis of randomized controlled trials. *Psychosom Med* 2014;76(3):190-196.

Shardell M, Hicks GE, Miller RR, Kritchevsky S, Andersen D, Bandinelli S, Cherubini A, Ferrucci L. Association of low vitamin D levels with the frailty syndrome in men and women. *J Gerontol A Biol Sci Med Sci* 2009;64(1):69-75.

Smit E, Crespo CJ, Michael Y, Ramirez-Marrero FA, Brodowicz GR, Bartlett S, Andersen RE. The effect of vitamin D and frailty on mortality among non-institutionalized US older adults. *Eur J Clin Nutr* 2012;66(9):1024-1028.

Smith RC, Gardiner JC, Lyles JS, Sirbu C, Dwamena FC, Hodges A, Collins C, Lein C, Given CW, Given B, Goddeeris J. Exploration of DSM-IV criteria in primary care patients with medically unexplained symptoms. *Psychosom Med* 2005;67(1):123-129.

Soysal P, Veronese N, Thompson T, Kahl KG, Fernandes BS, Prina AM, Solmi M, Schofield P, Koyanagi A, Tseng PT, Lin PY, Chu CS, Cosco TD, Cesari M, Carvalho AF, Stubbs B. Relationship between depression and frailty in older adults: a systematic review and meta-analysis. *Ageing Res Rev* 2017;36:78-87.

Spedding S. Vitamin D and depression: a systematic review and meta-analysis comparing studies with and without biological flaws. *Nutrients* 2014;6(4):1501-1518.

Stewart JC, Rand KL, Muldoon MF, Kamarck TW. A prospective evaluation of the directionality of the depression-inflammation relationship. *Brain Behav Immun* 2009;23(7):936-944.

Tiemeier H, Hofman A, van Tuijl HR, Kiliaan AJ, Meijer J, Breteler MM. Inflammatory proteins and depression in the elderly. *Epidemiology* 2003;14(1):103-107.

Thombs BD, Ziegelstein RC, Pilote L, Dozois DJ, Beck AT, Dobson KS, Fuss S, de Jonge P, Grace SL, Stewart DE, Ormel J, Abbey SE. Somatic symptom overlap in Beck depression inventory-II scores following myocardial infarction. *Br J Psychiatry* 2010;197(1):61-66.

Tyrka AR, Price LH, Kao HT, Porton B, Marsella SA, Carpenter LL. Childhood maltreatment and telomere shortening: preliminary support for an effect of early stress on cellular aging. *Biol Psychiatry* 2010;67:531-534.

Verhoeven JE, Révész D, Epel ES, Lin J, Wolkowitz OM, Penninx BW. Major depressive disorder and accelerated cellular aging: results from a large psychiatric cohort study. *Mol Psychiatry* 2014;19(8):895-901.

Wessely SS. Functional somatic syndromes: one or many? *Lancet* 1999;354(9182):936-939.

Wilhelm-Leen ER, Hall YN, Deboer IH, Chertow GM. Vitamin D deficiency and frailty in older Americans. *J Intern Med* 2010;268(2):171-80.

Wilkins CH, Mathews J, Sheline YI. Late life depression with cognitive impairment: Evaluation and treatment. *Clin Interv Aging* 2009;4:51-57.

Willeit P, Willeit J, Mayr A, Weger S, Oberhollenzer F, Brandstatter A, Kronenberg F, Kiechl S. Telomere length and risk of incident cancer and cancer mortality. JAMA 2010;304(1):69-75.

Wolkowitz OM, Reus VI, Mellon SH. Of sound mind and body: depression, disease, and accelerated aging. Dialogues Clin Neurosci 2011;13(1):25-39.

World Health Organisation (WHO). World report on ageing and health 2015 (www.who.int).

Zalli A, Jovanova O, Hoogendijk WJ, Tiemeier H, Carvalho LA. Low-grade inflammation predicts persistence of depressive symptoms. Psychopharmacology 2016;233(9):1669-1678.



Part I

Ageing-related biomarkers of physical frailty in late-life depression

Chapter 2

The relation between physical frailty and low-grade inflammation in late-life depression

MHL Arts, RM Collard, HC Comijs, PJW Naudé, R Risselada, P Naarding, RC Oude Voshaar
J Am Geriatr Soc 2015;63(8):1652-1657

Chapter 3

Telomere length and physical frailty in late-life depression

MHL Arts, RM Collard, HC Comijs, L de Jonge, P Naarding, B Penninx, Kok RM,
RC Oude Voshaar
Exp Gerontol 2018;111:229-234

Chapter 4

Vitamin D deficiency and course of frailty in a depressed older population

KS vd Berg, MHL Arts, RM Collard, RHS van den Brink, HC Comijs, RM Marijnissen,
RC Oude Voshaar
Aging Ment Health 2018;15:1-7

